



Oscillatory Behavior of Higher Order Nonlinear Difference Equations

Said R. Grace^a and John R. Graef^b

^a *Cairo University*

Department of Engineering Mathematics, Faculty of Engineering, Orman,
12221 Giza, Egypt

^b *University of Tennessee at Chattanooga*

Department of Mathematics, University of Tennessee at Chattanooga, TN
37403 Chattanooga, USA

E-mail(*corresp.*): John-Graef@utc.edu

E-mail: saidgrace@yahoo.com

Received November 4, 2019; revised June 28, 2020; accepted June 30, 2020

Abstract. The authors present some new oscillation criteria for higher order nonlinear difference equations with nonnegative real coefficients of the form

$$\Delta \left((a(t) (\Delta^{n-1} x(t))^\alpha) + q(t) x^\beta(t-m+1) = 0. \right.$$

Both of the cases n even and n odd are considered. They give examples to illustrate their results.

Keywords: oscillation, higher order, difference equations.

AMS Subject Classification: 34N05; 39A10; 34A21.

1 Introduction

In this paper, we study the oscillatory behavior of all solutions of the higher order nonlinear difference equation

$$\Delta \left((a(t) (\Delta^{n-1} x(t))^\alpha) + q(t) x^\beta(t-m+1) = 0, \quad t \geq t_0, \quad (1.1) \right.$$

where $n \geq 3$. We assume throughout that:

- (i) α and β are the ratios of odd positive integers with $\alpha \geq \beta$;

Copyright © 2020 The Author(s). Published by VGTU Press

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

(ii) $\{a(t)\}$ and $\{q(t)\}$ are sequences of positive real numbers for $t \geq t_0$;

(iii) $m > 1$.

We let

$$A(v, u) = \sum_{s=u}^v \frac{1}{a^{1/\alpha}(s)}, \quad v \geq u \geq t_0,$$

and assume that

$$\lim_{t \rightarrow \infty} A(t, t_0) < \infty. \quad (1.2)$$

By a *solution* of Equation (1.1), we mean a real sequence $\{x(t)\}$ defined for all $t \geq t_0 - m + 1$ that satisfies Equation (1.1) for all $t \geq t_0$. A solution of Equation (1.1) is called *oscillatory* if its terms are neither eventually positive nor eventually negative, and it is called *nonoscillatory* otherwise. If all solutions of the equation are oscillatory, then the equation itself is called oscillatory.

In recent years there has been much research activity concerning the oscillation and asymptotic behavior of solutions of various classes of difference equations and we mention [1, 2, 3, 4, 5, 6, 7] and the references cited therein as examples of some recent contributions in this area. There have been numerous studies on second order difference equations due to their use in the natural sciences and as well as for theoretical interests. Recent results on the oscillatory and asymptotic behavior of solutions of second order difference equations can be found, for example, in [8, 9, 10, 11, 12, 13, 17, 18, 19]. However, it appears that there are no known results regarding the oscillation of solutions of higher order difference equations of the form of Equation (1.1). In view of this, our aim in this paper is to present some new sufficient conditions that ensure that all solutions of Equation (1.1) are oscillatory.

2 Main results

We begin with some useful lemmas.

Lemma 1. ([1, Theorem 1.8.11], [3, Lemma 1]) *Let $\{x(t)\}$ be defined for $t \geq t_0$ and $x(t) > 0$ with $\Delta^n x(t)$ of constant sign and not identically zero for $t \geq t_0$. Then there exists an integer p , $0 \leq p \leq n$, with $p + n$ even if $\Delta^n x(t) \geq 0$ and $p + n$ odd for $\Delta^n x(t) \leq 0$, such that:*

i) $p \leq n - 1$ implies $(-1)^{p+i} \Delta^i x(t) > 0$ for $t \geq t_0$ and $p \leq i \leq n - 1$;

ii) $p \geq 1$ implies $\Delta^i x(t) > 0$ for all large $t \geq t_0$ and $1 \leq i \leq p - 1$.

Lemma 2. ([1, Corollary 1.8.12], [3, Lemma 2]) *Let $\{x(t)\}$ be defined for $t \geq t_0$ with $x(t) > 0$ and $\Delta^n x(t) \leq 0$ and not identically zero for $t \geq t_0$. Then there exists $t_1 \geq t_0$ such that*

$$x(t) \geq \frac{1}{(n-1)!} (t-t_1)^{n-1} \Delta^{n-1} x(2^{n-p-1}t) \quad \text{for } t \geq t_1,$$

where p is defined as in Lemma 1. Furthermore, if $\{x(t)\}$ is increasing, then

$$x(t) \geq \frac{1}{(n-1)!} \left(\frac{t}{2^{n-1}} \right)^{n-1} \Delta^{n-1} x(t) \quad \text{for } t \geq t_1.$$

The following lemma is an extension of the discrete analogue of known results in [15] and [16, Corollary 1]; it can also be found in [2, Lemma 6.2.2] and [14, Corollary 7.4.1]. The proof is immediate.

Lemma 3. *Let $\{q(t)\}$ be a sequence of positive real numbers, m and p be positive numbers, and $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous nondecreasing function with $xf(x) > 0$ for $x \neq 0$. If the first order delay inequality*

$$\Delta y(t) + q(t)f(y(t - m + 1)) \leq 0$$

has an eventually positive solution, then so does the delay equation

$$\Delta y(t) + q(t)f(y(t - m + 1)) = 0.$$

We are now ready for our first oscillation result; it is for the case where n is even.

Theorem 1. *Let n be even and assume that there is a number k such that $m > (n - 2)k + 1$. If the first order equations*

$$\begin{aligned} \Delta Y(t) + q(t) \left(\frac{1}{2^{n-1}(n-1)!} \right)^\beta & \left((t - m + 1)^{(n-1)\beta} a^{-\beta/\alpha}(t - m + 1) \right) \\ & \times Y^{\beta/\alpha}(t - m + 1) = 0, \end{aligned} \tag{2.1}$$

$$\begin{aligned} \Delta Z(t) + \left(\frac{1}{a(t)} \sum_{s=t_0}^{t-1} \left(\frac{1}{2^{n-2}(n-2)!} \right)^\beta q(s) (s - m + 1)^{(n-2)\beta} \right)^{1/\alpha} \\ & \times Z^{\beta/\alpha}(t - m + 1) = 0, \end{aligned} \tag{2.2}$$

and

$$\Delta W(t) + \frac{(k^{n-2})^{\beta/\alpha}}{a^{1/\alpha}(t)} \left(\sum_{s=t_0}^{t-1} q(s) \right)^{1/\alpha} W^{\beta/\alpha}(t - m + (n - 2)k + 1) = 0,$$

are oscillatory, then Equation (1.1) is oscillatory.

Proof. Let $\{x(t)\}$ be a nonoscillatory solution of Equation (1.1), say $x(t) > 0$ and $x(t - m + 1) > 0$ for $t \geq t_1$ for some $t_1 \geq t_0$. It follows from Equation (1.1) that

$$\Delta \left(a(t) (\Delta^{n-1}x(t))^\alpha \right) \leq -q(t)x^\beta(t - m + 1) \leq 0 \tag{2.3}$$

for $t \geq t_1$. Hence, $a(t) (\Delta^{n-1}x(t))^\alpha$ is nonincreasing and eventually of one sign. That is, there exists $t_2 \geq t_1$ such that

$$(I) \ \Delta^{n-1}x(t) > 0 \quad \text{or} \quad (II) \ \Delta^{n-1}x(t) < 0 \quad \text{and} \quad \Delta^{n-2}x(t) > 0 \quad \text{for} \quad t \geq t_2.$$

First, we consider Case (I). By Lemma 1, we have $\Delta x(t) > 0$ for $t \geq t_2$. From Lemma 2, we see that

$$x(t) \geq \frac{1}{(n-1)!} \left(\frac{t}{2^{n-1}} \right)^{n-1} \Delta^{n-1}x(t) \quad \text{for} \quad t \geq t_2,$$

and so

$$x(t - m + 1) \geq \frac{1}{(n - 1)!} \left(\frac{t - m + 1}{2^{n-1}} \right)^{n-1} \Delta^{n-1}x(t - m + 1) \text{ for } t \geq t_3, \tag{2.4}$$

for some $t_3 \geq t_2$. Using (2.4) in (2.3), we obtain

$$\begin{aligned} & \Delta \left(a(t) (\Delta^{n-1}x(t))^\alpha \right) \\ & \leq -q(t) \left(\frac{1}{2^{n-1}(n-1)!} \right)^\beta (t - m + 1)^{(n-1)\beta} (\Delta^{n-1}x(t - m + 1))^\beta \end{aligned}$$

or

$$\begin{aligned} \Delta Y(t) + q(t) \left(\frac{1}{2^{n-1}(n-1)!} \right)^\beta & \left((t - m + 1)^{(n-1)\beta} a^{-\beta/\alpha}(t - m + 1) \right) \\ & \times Y^{\beta/\alpha}(t - m + 1) \leq 0 \end{aligned} \tag{2.5}$$

for $t \geq t_3$, where $Y(t) = a(t) (\Delta^{n-1}x(t))^\alpha$. It follows from Lemma 3 that the corresponding Equation (2.1) also has a positive solution, which is a contradiction.

Next, we consider Case (II). Since n is even, we distinguish the following two cases (recall that (2.3) holds):

$$\text{(III) } x(t) > 0, \Delta x(t) > 0, \dots, \Delta^{n-2}x(t) > 0, \Delta^{n-1}x(t) < 0 \text{ for } t \geq t_2,$$

or

$$\text{(IV) } x(t) > 0, \Delta x(t) < 0, \dots, \Delta^{n-2}x(t) > 0, \Delta^{n-1}x(t) < 0 \text{ for } t \geq t_2.$$

If (III) holds, then by Lemma 2, we see that

$$x(t - m + 1) \geq \frac{1}{(n - 2)!} \left(\frac{t - m + 1}{2^{n-2}} \right)^{n-2} \Delta^{n-2}x(t - m + 1) \text{ for } t \geq t_3 \tag{2.6}$$

for some $t_3 \geq t_2$. Using (2.6) in (2.3), we have

$$\begin{aligned} & \Delta \left(a(t) (\Delta^{n-1}x(t))^\alpha \right) \\ & \leq -q(t) \left(\frac{1}{2^{n-2}(n-2)!} \right)^\beta (t - m + 1)^{(n-2)\beta} (\Delta^{n-2}x(t - m + 1))^\beta \end{aligned}$$

for $t \geq t_3$. Summing this inequality from t_3 to $t - 1$ gives

$$\begin{aligned} & - \left(a(t) (\Delta^{n-1}x(t))^\alpha \right) \\ & \geq \left(\sum_{s=t_3}^{t-1} \left(\frac{1}{2^{n-2}(n-2)!} \right)^\beta q(s) (s - m + 1)^{(n-2)\beta} \right) (\Delta^{n-2}x(t - m + 1))^\beta \end{aligned}$$

or

$$-\Delta^{n-1}x(t) \geq \left(\frac{1}{a(t)} \sum_{s=t_3}^{t-1} \left(\frac{1}{2^{n-2}(n-2)!} \right)^\beta q(s) (s-m+1)^{(n-2)\beta} \right)^{1/\alpha} \times (\Delta^{n-2}x(t-m+1))^{\beta/\alpha},$$

so

$$\Delta Z(t) + \left(\frac{1}{a(t)} \sum_{s=t_3}^{t-1} \left(\frac{1}{2^{n-2}(n-2)!} \right)^\beta q(s) (s-m+1)^{(n-2)\beta} \right)^{1/\alpha} \times Z^{\beta/\alpha}(t-m+1) \leq 0$$

for $t \geq t_3$, where $Z(t) = \Delta^{n-2}x(t) > 0$. The remainder of the proof in this case is similar to that of Case (I) and is omitted.

Finally, we consider Case (IV). We have

$$(-1)^i \Delta^i x(t) > 0 \text{ for } i = 0, 1, \dots, n-1, \text{ and } t \geq t_2. \tag{2.7}$$

Notice that

$$-\Delta^{n-3}x(t) \geq \Delta^{n-3}x(t+k) - \Delta^{n-3}x(t) = \sum_{s=t}^{t+k-1} \Delta^{n-2}x(s) \geq k(\Delta^{n-2}x(t+k)).$$

Repeating this $n-2$ times, we arrive at

$$x(t) \geq k^{n-2} \Delta^{n-2}(t + (n-2)k),$$

so

$$x(t-m+1) \geq k^{n-2} \Delta^{n-2}(t-m+(n-2)k+1).$$

Using this inequality in (2.3), we have

$$\Delta \left(a(t) (-\Delta^{n-1}x(t))^\alpha \right) \geq q(t) \left(k^{n-2} \Delta^{n-2}(t-m+(n-2)k+1) \right)^\beta.$$

Summing from t_2 to $t-1$ yields

$$a(t) (-\Delta^{n-1}x(t))^\alpha - a(t_2) (-\Delta^{n-1}x(t_2))^\alpha \geq \sum_{s=t_2}^{t-1} q(s) \left(k^{n-2} \Delta^{n-2}x(s-m+(n-2)k+1) \right)^\beta$$

or

$$-\Delta^{n-1}x(t) \geq \frac{(k^{n-2})^{\beta/\alpha}}{a^{1/\alpha}(t)} \left(\sum_{s=t_2}^{t-1} q(s) \right)^{1/\alpha} (\Delta^{n-2}x(t-m+(n-2)k+1))^{\beta/\alpha},$$

and hence we obtain

$$\Delta W(t) + \frac{(k^{n-2})^{\beta/\alpha}}{a^{1/\alpha}(t)} \left(\sum_{s=t_1}^{t-1} q(s) \right)^{1/\alpha} W^{\beta/\alpha}(t-m+(n-2)k+1) \leq 0,$$

where $W(t) = \Delta^{n-2}x(t) > 0$. The rest of the proof is similar to that of Case (I) and is left to the reader. This completes the proof of the theorem. \square

To obtain some consequences of the above theorem, we let

$$P(t) \leq \min \left\{ q(t) \left(\frac{1}{2^{n-1}(n-1)!} \right)^\beta (t-m+1)^{(n-1)\beta} a^{-\beta/\alpha} (t-m+1), \left(\frac{1}{a(t)} \right) \times \sum_{s=t_0}^{t-1} \left(\frac{1}{2^{n-2}(n-2)!} \right)^\beta q(s) (s-m+1)^{(n-2)\beta} \right\}^{\frac{1}{\alpha}}, \left(\frac{k^{n-2}}{a^{1/\alpha}(t)} \right)^{\frac{\beta}{\alpha}} \left(\sum_{s=t_0}^{t-1} q(s) \right)^{\frac{1}{\alpha}} \right\}.$$

and $\tau = \max\{m - (n - 2)k - 1, m - 1\}$.

The following corollaries are immediate consequences of known results.

Corollary 1. Let n be even and k be a number such that $m > (n - 2)k + 1$. If the first order delay equation

$$\Delta y(t) + P(t)y^{\beta/\alpha}(t - \tau) = 0$$

is oscillatory, then Equation (1.1) is oscillatory.

Corollary 2. Let n be even and k be a number such that $m > (n - 2)k + 1$. If

$$\liminf_{t \rightarrow \infty} \sum_{s=t-m+1}^{t-1} P(s) \begin{cases} > \frac{(\tau-1)^{\tau-1}}{\tau^\tau}, & \text{if } \beta = \alpha, \\ = \infty, & \text{if } \beta < \alpha, \end{cases}$$

then Equation (1.1) is oscillatory.

We now turn our attention to the case where n is odd.

Theorem 2. Let n be odd and k be a number such that $m > (n - 1)k + 1$. If the first order Equations (2.1)–(2.2), and

$$\Delta W(t) + q(t) \left(\frac{k^{n-1}}{a^{1/\alpha}(t)} \right)^\beta W^{\beta/\alpha}(t - m + (n - 1)k + 1) = 0 \tag{2.8}$$

are oscillatory, then Equation (1.1) is oscillatory.

Proof. Let $\{x(t)\}$ be a nonoscillatory solution of Equation (1.1), say $x(t) > 0$ and $x(t - m + 1) > 0$ for $t \geq t_1$ for some $t_1 \geq t_0$. As in the proof of Theorem 1, we see that $a(t) (\Delta^{n-1}x(t))^\alpha$ is nonincreasing and eventually of one sign. That is, there exists $t_2 \geq t_1$ such that

$$(I) \Delta^{n-1}x(t) > 0 \quad \text{or} \quad (II) \Delta^{n-1}x(t) < 0 \text{ and } \Delta^{n-2}x(t) > 0 \text{ for } t \geq t_2.$$

If Case (I) holds, either we have (A) $\Delta x(t) > 0$, or (B) $\Delta x(t) < 0$ for $t \geq t_2$. By Lemma 2 and (A), we obtain (2.4) and (2.5) which leads to a contradiction. If (B) holds, we see that (2.7) holds, and as in the proof of Theorem 1, we have

$$x(t - m + 1) \geq k^{n-1} \Delta^{n-1}(t - m + (n - 1)k + 1).$$

Thus,

$$\Delta \left(a(t) (\Delta^{n-1} x(t))^\alpha \right) \leq -q(t) (k^{n-1} \Delta^{n-1} x(t - m + (n - 1)k + 1))^\beta$$

or

$$\Delta W(t) \leq -q(t) \left(\frac{k^{n-1}}{a^{1/\alpha}(t - m + (n - 1)k + 1)} \right)^\beta W^{\beta/\alpha}(t - m + (n - 1)k + 1),$$

where $W(t) = a(t) (\Delta^{n-1} x(t))^\alpha > 0$. The rest of the proof is similar to that of the proof of Case I in Theorem 1 and hence is omitted. \square

Next, we consider Case (II). By Lemma 2 and as in the proof of Theorem 1, we obtain (2.6). The rest of the proof is similar to that of Case (III) in Theorem 1 and so we omit the details. This proves the theorem.

Now let

$$P^*(t) \leq \min \left\{ q(t) \left(\frac{1}{2^{n-1}(n-1)!} \right)^\beta \left((t-m+1)^{(n-1)\beta} a^{-\beta/\alpha}(t-m+1) \right), \right. \\ \left. \left(\frac{1}{a(t)} \sum_{s=t_0}^{t-1} \left(\frac{1}{2^{n-2}(n-2)!} \right)^\beta q(s) (s-m+1)^{(n-2)\beta} \right)^{\frac{1}{\alpha}}, \frac{(k^{n-1})^{\frac{\beta}{\alpha}}}{a^{1/\alpha}(t)} \left(\sum_{s=t_0}^{t-1} q(s) \right)^{\frac{1}{\alpha}} \right\}$$

and $\tau^* = \max\{m - (n - 1)k - 1, m - 1\}$.

The following corollaries are analogous to those for the case where n is even.

Corollary 3. Let n be odd and k be a number such that $m > (n - 1)k + 1$. If the first order delay equation

$$\Delta y(t) + P^*(t)y^{\beta/\alpha}(t - \tau^*) = 0$$

is oscillatory, then Equation (1.1) is oscillatory.

Corollary 4. Let n be odd and k be a number with $m > (n - 1)k + 1$. If

$$\liminf_{t \rightarrow \infty} \sum_{s=t-m+1}^{t-1} P^*(s) \begin{cases} > \frac{(\tau^* - 1)^{\tau^*-1}}{\tau^* \tau^*}, & \text{if } \beta = \alpha, \\ = \infty, & \text{if } \beta < \alpha, \end{cases}$$

then Equation (1.1) is oscillatory.

Remark 1. Clearly our results hold if $n = 2$, and they appear to be new in this case as well.

If we replace condition (1.2) by

$$\lim_{t \rightarrow \infty} A(t, t_0) = \infty, \tag{2.9}$$

it is easy to obtain the following results.

Theorem 3. Let n be even, condition (2.9) hold, and there exists a number k with $m > (n - 2)k + 1$. If the first order Equation (2.1) is oscillatory, then Equation (1.1) is oscillatory.

Theorem 4. Let n be odd, condition (2.9) hold, and k be a number with $m > (n - 1)k + 1$. If the first order Equations (2.1) and (2.8) are oscillatory, then Equation (1.1) is oscillatory.

As an example to illustrate our results, consider the difference equation

$$\Delta \left(\frac{1}{t^6} (\Delta^{n-1} x(t))^3 \right) + x^3(t - m + 1) = 0. \quad (2.10)$$

It is easy to see that if n is even and $m > (n - 2)k + 1$ (n is odd and $m > (n - 1)k + 1$) the conditions of Corollary 2.2 (respectively Corollary 2.4) are satisfied and hence we conclude that Equation (2.10) is oscillatory.

We conclude this paper with a suggestion for future research, namely, to study the oscillatory behavior of solutions of Equation (1.1) in case $\beta > \alpha$.

References

- [1] R.P. Agarwal. *Difference Equations and Inequalities: Theory, Methods, and Applications, 2nd Ed.* Monographs and Textbooks in Pure and Applied Mathematics, Dekker, New York, 2000. <https://doi.org/10.1201/9781420027020>.
- [2] R.P. Agarwal, M. Bohner, S.R. Grace and D. O'Regan. *Discrete Oscillation Theory.* Hindawi, New York, 2005. <https://doi.org/10.1155/9789775945198>.
- [3] R.P. Agarwal and S.R. Grace. Oscillation of higher order difference equations. *Appl. Math. Letters*, **13**:81–88, 2000. [https://doi.org/10.1016/S0893-9659\(99\)00149-4](https://doi.org/10.1016/S0893-9659(99)00149-4).
- [4] R.P. Agarwal, S.R. Grace and D. O'Regan. *Oscillation Theory for Difference and Functional Differential Equations.* Kluwer, Dordrecht, 2000. <https://doi.org/10.1007/978-94-015-9401-1>.
- [5] C. Dharuman, J.R. Graef, E. Thandapani and K.S. Vidhyaa. Oscillation of second order difference equations with a sub-linear neutral term. *J. Math. Appl.*, **40**:59–67, 2017. <https://doi.org/10.7862/rf.2017.4>.
- [6] H.A. El-Morshedy. Oscillation and nonoscillation criteria for half-linear second order difference equations. *Dynam. Systems Appl.*, **15**:429–450, 2006.
- [7] H.A. El-Morshedy. New oscillation criteria for second order linear difference equations with positive and negative coefficients. *Comput. Math. Appl.*, **58**:1988–1997, 2009. <https://doi.org/10.1016/j.camwa.2009.07.078>.
- [8] H.A. El-Morshedy and S.R. Grace. Comparison theorems for second order nonlinear difference equations. *J. Math. Anal. Appl.*, **306**:106–121, 2005. <https://doi.org/10.1016/j.jmaa.2004.12.024>.
- [9] S.R. Grace, R.P. Agarwal, M. Bohner and D. O'Regan. Oscillation of second-order strongly superlinear and strongly sublinear dynamic equations. *Commun. Nonlinear Sci. Numer. Simul.*, **14**:3463–3471, 2009. <https://doi.org/10.1016/j.cnsns.2009.01.003>.

- [10] S.R. Grace, M. Bohner and R.P. Agarwal. On the oscillation of second-order half-linear dynamic equations. *J. Difference Equ. Appl.*, **15**:451–460, 2009. <https://doi.org/10.1080/10236190802125371>.
- [11] S.R. Grace and H.A. El-Morshedy. Oscillation criteria of comparison type for second order difference equations. *J. Appl. Anal.*, **6**:87–103, 2000. <https://doi.org/10.1515/JAA.2000.87>.
- [12] S.R. Grace and J.R. Graef. Oscillatory behavior of second order nonlinear differential equations with a sublinear neutral term. *Math. Model. Anal.*, **23**:217–226, 2018. <https://doi.org/10.3846/mma.2018.014>.
- [13] J.R. Graef, S.R. Grace and E. Tunç. Oscillatory behavior of even-order nonlinear differential equations with a sublinear neutral term. *Opuscula Math.*, **39**:39–47, 2019. <https://doi.org/10.7494/OpMath.2019.39.1.39>.
- [14] I. Györi and G. Ladas. *Oscillation Theory of Delay Differential Equations and Applications*. Oxford University Press, Oxford, 1991.
- [15] G. Ladas and I.P. Stavroulakis. Oscillation caused by several retarded and advanced arguments. *J. Differ. Equations*, **44**:134–152, 1982. [https://doi.org/10.1016/0022-0396\(82\)90029-8](https://doi.org/10.1016/0022-0396(82)90029-8).
- [16] Ch.G. Philos. On the existence of nonoscillatory solutions tending to zero at ∞ for differential equations with positive delays. *Arch. Math. (Basel)*, **36**:168–178, 1981. <https://doi.org/10.1007/BF01223686>.
- [17] S. Selvarangam, M. Madhan and E. Thandapani. Oscillation theorems for second order nonlinear neutral type difference equations with positive and negative coefficients. *Rom. J. Math. Comput. Sci.*, **7**:1–10, 2017.
- [18] S. Selvarangam, E. Thandapani and S. Pinelas. Oscillation theorems for second order nonlinear neutral difference equations. *J. Inequal. Appl.*, **2014**(417):1–15, 2014. <https://doi.org/10.1186/1029-242X-2014-417>.
- [19] M.K. Yildiz and H. Ögünmez. Oscillation results of higher order nonlinear neutral delay difference equations with a nonlinear neutral term. *Hacet. J. Math. Stat.*, **43**:809–814, 2014.